


## GEOMETRY (FRACTAL AND OTHERWISE)

1. A well-known geometry theorem states that the sum of the interior angles of a triangle must be  $180^\circ$ . This problem guides you through finding a similar formula for the sum of the interior angles of any convex polygon. Convex means that every interior angle is less than  $180^\circ$ , so the first quadrilateral below is convex, but the second is not.<sup>1</sup>



a) Determine the sum of the interior angles of a convex quadrilateral (hint: divide it into two triangles).

b) Determine the sum of the interior angles of a convex pentagon (hint: divide it into a triangle and a quadrilateral, then use your formula from part a).

c) Find a general formula for the sum of interior angles of a convex polygon with  $n \geq 3$  sides, then prove that your formula always works.

---

*Date:* December 8, 2021.

<sup>1</sup>Technically, convex means that the straight line between any two points of the polygon is entirely inside the polygon

**Definition.** We construct the Cantor middle-thirds set as follows. Define the following sets (as unions of intervals in  $\mathbb{R}$ ):

$$\begin{aligned} C_0 &= [0, 1] \\ C_1 &= \left[0, \frac{1}{3}\right] \cup \left[\frac{2}{3}, 1\right] \\ C_2 &= \left[0, \frac{1}{9}\right] \cup \left[\frac{2}{9}, \frac{1}{3}\right] \cup \left[\frac{2}{3}, \frac{7}{9}\right] \cup \left[\frac{8}{9}, 1\right] \\ &\vdots \end{aligned}$$

In general,  $C_{k+1}$  is formed by removing the open middle third of each interval in  $C_k$ . The **Cantor middle-thirds set** is

$$\mathcal{C} = \bigcap_{k=0}^{\infty} C_k$$

2. Prove or disprove the following statements about  $\mathcal{C}$ .

a)  $\mathcal{C} = \emptyset$

b)  $\forall n \in \mathbb{N}, \left(0, \frac{1}{2^n}\right) \not\subseteq \mathcal{C}$

**Definition.** Any real number can be expressed in ternary, which is like binary but with 3 possible values for each digit. For example, the ternary number 201 is

$$(2 \times 3^2) + (0 \times 3^1) + (1 \times 3^0) = 18 + 0 + 1 = 19.$$

And the ternary number 0.201 is

$$(2 \times 3^{-1}) + (0 \times 3^{-2}) + (1 \times 3^{-3}) = \frac{2}{3} + \frac{0}{9} + \frac{1}{27} = \frac{19}{27}.$$

**Proposition 1.** *Let  $x \in [0, 1]$ . Then  $x \in \mathcal{C}$  if and only if  $x$  can be written as a ternary number using **only** the digits 0 and 2.*

3. Is  $1/4$  in the Cantor set?

4. Is the Cantor set countable?

**Challenge.** Prove proposition 1.