

EXAM 2 REVIEW

Definition. A **random variable** X assigns a number to each outcome in the sample space S .

- (1) All random variables have a **cumulative distribution function (CDF)**: $F(x) = P(X \leq x)$.
- (2) A discrete random variable has a **probability mass function (PMF)**: $p(x) = P(X = x)$.
- (3) A continuous random variable has a **probability density function (PDF)** $f(x)$ such that for any numbers a and b (with $a \leq b$)

$$P(a \leq X \leq b) = \int_a^b f(x)dx$$

Definition. The **expected value** (or **mean**, often denoted by μ):

- (1) If X is a discrete RV with PMF $p(x)$, then $E(X) = \sum_x xp(x)$.
- (2) If X is a continuous RV with PDF $f(x)$, then $E(X) = \int_{-\infty}^{\infty} xf(x)dx$.

Definition. The **variance** (often denoted by σ^2): $\text{Var}(X) = E[(X - \mu)^2]$. The **standard deviation**: $\sigma = \sqrt{\text{Var}(x)}$.

Theorem. For any random variable X and any constants a and b :

- (1) $E(aX + b) = aE(X) + b$ and
- (2) $\text{Var}(aX + b) = a^2 \text{Var}(X)$.

Theorem. If X_1, X_2, \dots, X_n are independent, then

- (1) $E(X_1 + X_2 + \dots + X_n) = E(X_1) + E(X_2) + \dots + E(X_n)$
- (2) $\text{Var}(X_1 + X_2 + \dots + X_n) = \text{Var}(X_1) + \text{Var}(X_2) + \dots + \text{Var}(X_n)$

Definition. A **random sample of size n** is a set of independent identically distributed random variables X_1, X_2, \dots, X_n . Some **sample statistics**:

- (1) The **sample mean**: $\bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$
- (2) The **sample variance**: $S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$

Theorem. For any random sample from a population with mean μ and variance σ^2 :

- (1) $E(\bar{X}) = \mu$ and $\text{Var}(\bar{X}) = \frac{\sigma^2}{n}$
- (2) $E(S^2) = \sigma^2$

Definition. The **sample standard error** is $\frac{s}{\sqrt{n}}$

Definition. A sample statistic \hat{X} is an **unbiased estimator** of population parameter ρ if $E(\hat{X}) = \rho$.

Theorem. If \bar{X} is the mean of a random sample from a normally distribution population, then \bar{X} is normally distributed (with mean and variance given in the last theorem).

Theorem (Central Limit Theorem). If \bar{X} is the mean of a random sample from a population, then \bar{X} is approximately normally distributed (with mean and variance given in the theorem above).

A. CONFIDENCE (AND PREDICTION) INTERVALS

1. $100(1 - \alpha)\%$ CI for μ (known σ): $\bar{x} \pm z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$

2. $100(1 - \alpha)\%$ CI for μ (large sample): $\bar{x} \pm z_{\alpha/2} \frac{s}{\sqrt{n}}$

3. $100(1 - \alpha)\%$ CI for μ (normal population): $\bar{x} \pm t_{\alpha/2, n-1} \frac{s}{\sqrt{n}}$

4. $100(1 - \alpha)\%$ **prediction interval** for μ (normal population): $\bar{x} \pm t_{\alpha/2, n-1} \sqrt{\frac{s^2(n+1)}{n}}$

5. $100(1 - \alpha)\%$ CI for $\mu_1 - \mu_2$ (known σ_1 and σ_2 , normal populations): $\bar{x} - \bar{y} \pm z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}$

6. $100(1 - \alpha)\%$ CI for $\mu_1 - \mu_2$ (large samples): $\bar{x} - \bar{y} \pm z_{\alpha/2} \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$

7. $100(1 - \alpha)\%$ CI for $\mu_1 - \mu_2$ (normal populations with the same variance):

$$\bar{x} - \bar{y} \pm t_{\alpha/2, n_1+n_2-2} \sqrt{s_p^2 \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}$$

$$s_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2} \text{ is the pooled estimator of the common variance}$$

8. $100(1 - \alpha)\%$ CI for $\mu_1 - \mu_2$ (normal populations with difference variances): $\bar{x} - \bar{y} \pm t_{\alpha/2, \nu} \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$

$$\nu \approx \frac{\left(\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)^2}{\frac{\left(\frac{s_1^2}{n_1} \right)^2}{n_1 - 1} + \frac{\left(\frac{s_2^2}{n_2} \right)^2}{n_2 - 1}} \text{ (round down to the nearest integer)}$$

9. $100(1 - \alpha)\%$ CI for a proportion θ (large sample; x and $n - x$ both large): $\frac{x}{n} \pm z_{\alpha/2} \sqrt{\frac{1}{n} \left(\frac{x}{n} \right) \left(1 - \frac{x}{n} \right)}$

B. PRACTICE PROBLEMS

1. I have collected the heights of 71 adult humans, more or less at random. This sample has mean $\bar{x} = 68.28$ and standard deviation of $s = 3.60$. Calculate a 90% confidence interval for the mean height of an adult human.
2. A January, 2016 YouGov poll of 388 likely South Carolina Democratic primary voters found that 60% would vote for Clinton. Calculate a 95% confidence interval for the true percentage of Clinton voters in the South Carolina Democratic primary (which was held on Saturday, February 27 and in which Clinton got 73.5% of the votes).
3. The article “Orchard Floor Management Utilizing Soil-Applied Coal Dust for Frost Protection” reports the following values for soil heat flux of $n = 8$ plots covered with coal dust:

34.7 35.4 34.7 37.7 32.5 28.0 18.4 24.9

The mean soil heat flux for plots covered only with grass is 29.0. Assuming that soil heat flux is normally distributed, does this data evidence that coal dust is effective in increasing mean soil heat flux? Use a confidence interval to justify your conclusion.

4. It is thought that roughly $\frac{2}{3}$ of all people have a dominant right foot and $\frac{2}{3}$ have a dominant right eye. Do people also kiss to the right? The article “Human Behavior: Adult Persistence of Head-Turning Asymmetry” reported that in a random sample of 124 kissing couples, 80 of the couples tended to lean more to the right than left. Does this result suggest that more than half of all couples lean right when kissing? Does this result provide evidence against the hypothesis that $\frac{2}{3}$ of all kissing couples lean right? Use confidence intervals to support your conclusions.
5. Suppose that a population is uniformly distributed on the interval $[1, \beta]$. Let X_1, X_2, \dots, X_n be a random sample. Find an unbiased estimator for β .

Challenge. Continue with the situation of the previous problem. Find another unbiased estimator for β .

Challenge. Sometimes people conducting surveys want to ask questions with potentially embarrassing answers. They may use something called a *randomized response technique* to make sure that survey participants have no reason to lie. For example, suppose you wanted to determine how widespread cheating is on campus. You could print the phrase “I have cheated on a test” on 4 cards and “I have never cheated on a test” on 6 cards. Then you could ask each survey participant to select a random card and tell you if the statement on the card is true or false *without showing you the card*.

- a) Suppose 55% of those surveyed said the statement on the card was true. What is your estimate of the percentage of Gonzaga students who have cheated?
- b) Suppose 60% of those surveyed said the statement on the card was true. What is your estimate of the percentage of Gonzaga students who have cheated?
- c) Suppose 65% of those surveyed said the statement on the card was true. What is your conclusion?